

Research statement

The results obtained during the work on my thesis and several years after its defense are in the field of model theory, and its applications to algebraic geometry. More recently I have taken interest in non-Archimedean geometry and motivic integration. I have been inspired to start working in this field (and in particular on Project 2a below) after learning from a conversation with E. Hrushovski about a somewhat overlooked tool, the asymptotic valued field, defined by Abraham Robinson using ultraproducts of real closed fields, that allows one to relate collapsed Gromov-Hausdorff limits to geometry of non-Archimedean analytic spaces.

I will now quickly summarize my work in model theory and then proceed to describing my current research projects.

1. In a joint paper with Assaf Hasson [HS17] we have settled Restricted Trichotomy conjecture of Boris Zilber in dimension 1. This conjecture plays a crucial role in Zilber's model-theoretic proof [Zil14] of a strengthening of Bogomolov's Torelli-type theorem for curves over finite fields [BKT10]. Our theorem can be regarded as an analogue of the Fundamental Theorem of abstract projective geometry [Art57, Har67].
2. In papers [Sus16] and [SSZ14] (joint work with Boris Zilber and Vinesh Solanki) I have studied certain classes of model-theoretic structures called Zariski geometries and developed a criterion that they can be represented using Zariski closed subsets of Cartesian powers of algebraic varieties.

Project 1. Refined tropical multiplicities and motivic integraton.

Let Δ be a lattice polygon, and consider the projective toric surface $Y(\Delta)$ associated to it. Let $|L(\Delta)|$ be the linear system containing all curves with Newton polygon Δ . The *toric Severi degree* $n^{\Delta, \delta}$ is the number of integral curves in the linear system $|L(\Delta)|$ that have δ nodes and that pass through $n - \delta$ points in general position, where n is the dimension of the linear system $|L(\Delta)|$. Denote by g the genus of the general smooth element of $L(\Delta)$. According to an idea proposed by Yau and Zaslow, and developed by Beauville, Fantechi, Göttsche and van Straten [Bea99], [FGvS99], if $C \rightarrow T$ is a family of curves on a smooth surface where only finitely many rational curves occur, and if these curves have only nodal singularities while the rest of the fibres are smooth, the number of curves with g nodes can be computed as the Euler characteristic of the relative compactified Jacobian $\text{Jac}(C/T)$ of C over T .

This technique can be generalized to the case of strictly positive cogenus $g - \delta$; provided that the family C has finitely many curves with δ nodes and all other fibres are of geometric genus greater than $g - \delta$ one can define numbers n_1, \dots, n_δ by performing variable change on the generating series of Euler characteristics of relative Hilbert schemes of i points on C/T , and $n^{\Delta, \delta} = n_\delta$ [KST11], [PT10]. In [GS14] Göttsche and Shende have proposed to replace the Euler characteristic with Hirzebruch's χ_y -genus in this construction. They show that the generating series mentioned above give rise to a well-defined *refined count* $N_\delta \in \mathbb{Z}[y, y^{-1}]$.

Both toric Severi degrees and refined toric Severi degrees can be computed tropically as a sum over tropical curves $\Gamma \subset \mathbb{R}^2$ of degree Δ and genus $g - \delta$ passing through $n - \delta$ points in general position of (refined) *tropical multiplicities* $N(\Gamma)$ of Γ , the latter is computed combinatorially in terms of the shape of the graph Γ . In [NPS16] Nicaise, Payne and Schroeter propose a geometric interpretation of the refined tropical multiplicities of Block and Göttsche using the motivic volume of Hrushovski and Kazhdan [HK06]. Motivic volume is a map $\text{Vol} : K_0(\text{VF}) \rightarrow K_0(\text{Var}_k)$ from the Grothendieck ring of non-Archimedean semi-algebraic subsets of algebraic varieties defined over an algebraically closed valued field to the Grothendieck ring of varieties over the residue field, and is characterized by the following two properties:

1. if $\Delta \subset \mathbb{R}^n$ is a polytope $\text{Vol}(\text{trop}^{-1}(\Delta)) = \chi'(\Delta)(\mathbb{L} - 1)^{\dim \Delta}$ where \mathbb{L} is the class of the affine line, χ' is the modified Euler characteristic $\chi'(\Delta) = \lim_{a \rightarrow \infty} \chi_c(X \cap [-a, a]^n)$ and χ_c is the Euler characteristic with compact supports;
2. let \mathcal{X} be flat projective scheme over the value ring R such that $\mathcal{X} \otimes_R K \cong X$ and let $\text{sp} : X(K) \rightarrow \mathcal{X}_0$ be the specialization map. If $U \subset \mathcal{X}_0$ is a constructible subset then $\text{Vol}(\text{sp}^{-1}(U)) = [U]$.

Additive invariants of varieties, such as Euler characteristic and χ_y -genus, can be lifted to $K_0(\text{VF})$ by evaluating them on the motivic volume of the semi-algebraic set. Let S be a set of $n - \delta$ points on the dense torus in $Y(\Delta)$ in general position. If $\Gamma \subset \mathbb{R}^2$ is a tropical curve then denote $C_\Gamma \rightarrow T_\Gamma$ the universal family of curves on \mathbb{P}^2 of degree Δ and passing through S that tropicalize to Γ . Nicaise, Payne and Schroeter make the following Conjectures [NPS16] that they check for $g = 1$:

Conjecture 1. Assume that $\delta = g$ and let $\Gamma \subset \mathbb{R}^2$ be a rational tropical curve of degree Δ passing through points in $\text{trop}(S)$. Then the refined tropical multiplicity $N(\Gamma)$ is equal to $y^{-g}\chi_y(\text{Vol}(\text{Jac}(C_\Gamma/T_\Gamma)))$.

Conjecture 2. For any integer $\delta \in [0, g]$, let Γ be a tropical curve in \mathbb{R}^2 of genus $g - \delta$ and degree Δ passing through $\text{trop}(S)$. Then refined tropical multiplicity $N(\Gamma)$ is equal to $y^{-\delta}N_\delta(C_\Gamma)$ where $N_\delta(C_\Gamma)$ is computed from the generating series similarly to $N_\delta(C)$.

In view of Conjecture 1 it is natural to try to compute the motivic volume of semi-algebraic families of Jacobians of smooth projective curves. According to computations of Nicaise, Payne and Schroeter, the χ_y -genus of the motivic volume of a semi-algebraic family of varieties can be non-zero even if the χ_y -genus of each fiber vanishes (as is the case for Abelian varieties, for example). Nevertheless, I have found the following

Theorem (S.[Sus18b]) The motivic volume of a semi-algebraic family of Abelian varieties that admits a fibrewise uniformization by an algebraic torus vanishes.

My next steps in this project are as follows. Firstly, I intend to prove an analogue of theorem above for relative Hilbert schemes of points of Mumford curves (that is, curves that admit a rigid-analytic uniformization). Secondly, I would like to find a formula for the χ_y -genus of a semi-algebraic subfamily of the relative Jacobian of a one-dimensional family of curves in terms of its monodromy, and similarly for semi-algebraic subfamilies of the relative Hilbert schemes of points. Finally, I want to study the relation between the χ_y -genus of such semi-algebraic sets and the tropical multiplicity $N(\Gamma)$.

Project 2. Degenerations of complex manifolds and non-Archimedean geometry

2a. Non-Archimedean limits of differential forms, Gromov-Hausdorff limits and SYZ conjecture.

In the beginning of the 2000s Kontsevich and Soibelman have come up with a new approach to Ströminger-Yau-Zaslow (SYZ) conjecture from the subject of mirror symmetry [KS04]. In a nutshell, this conjecture posits that two mirror partners X and X^\vee admit two fibrations over the same base B , a manifold of real dimension $1/2 \dim_{\mathbb{R}} X$, and that generic fibres of the two fibrations are tori that are dual to each other. Kontsevich and Soibelman suggest to look for the base of the SYZ fibration using two methods, non-Archimedean and differential-geometric.

Let $\mathcal{X} \rightarrow \mathbb{C}^\times$ be a family of projective complex manifolds with maximal unipotent monodromy at 0 and let \mathcal{L} be a relatively ample line bundle on $\mathcal{X}/\mathbb{C}^\times$. Let $X = \mathcal{X} \otimes \mathbb{C}((t))$ with respect to the natural inclusion $\mathbb{C}[t] \hookrightarrow \mathbb{C}((t))$ and let L be the restriction of \mathcal{L} to X . Denote \mathcal{X}_s the fibre over a point $s \in \mathbb{C}^\times$ and assume that $\{\omega_s\}_{s \in \mathbb{C}^\times}$ is a smooth family of Kähler forms of Ricci-flat metrics on \mathcal{X}_s for s in a punctured neighbourhood of 0, such that $\omega_s \in c_1(\mathcal{L}_s)$. Let $\tilde{\omega}_s$ be the Kähler form of the metric normalized so that the diameter of \mathcal{X}_s is 1.

On the one hand the base of the SYZ fibration is to be identified with a subset, called *essential skeleton* $\text{Sk}(X)$, of the Berkovich analytification X^{an} of the variety X , and on the other hand it is to be constructed as a Gromov-Hausdorff limit B of $(\mathcal{X}_s, \tilde{\omega}_s)$ as $s \rightarrow 0$ (“large complex structure limit”).

Conjecture A (Kontsevich and Soibelman, Gross and Wilson, Todorov) The limit space B contains an open dense subset B^{sm} which has the structure of an affine manifold, $\dim_{\mathbb{R}} B^{\text{sm}} = 1/2 \dim_{\mathbb{R}} \mathcal{X}_s$, and $B \setminus B^{\text{sm}}$ is of Hausdorff codimension 2. Moreover, B^{sm} is a Riemannian manifold such that the metric tensor is locally a Hessian (in the affine coordinates) of a potential function, $g_{ij} = \partial^2 F / \partial x_i \partial x_j$, and the potential satisfies the real Monge-Ampere equation $\det(\partial^2 F / \partial x_i \partial x_j) \equiv \text{const}$.

Conjecture B (Kontsevich and Soibelman) The non-Archimedean space X admits a continuous surjective map to $\text{Sk}(X)$ with the fibre non-Archimedean torus over an open dense subset of $\text{Sk}(X)$ with complement of codimension 2; this fibration induces a singular affine structure on $\text{Sk}(X)$.

Conjecture C (Konstevich and Soibelman) The two spaces are naturally isomorphic as (singular) affine manifolds.

In [Sus18a] I have considered the following toy version of Conjecture C: I consider maximally unipotent degenerations of complex curves \mathcal{X}_s of genus ≥ 1 endowed with flat (pseudo-)Kähler metrics with the Kähler form $\Omega_s \wedge \bar{\Omega}_s$ where Ω is a relative holomorphic 1-form.

Let \mathcal{X} be a model of X , i.e. a flat R -scheme such that $\mathcal{X} \otimes_R K \cong X$. Assume that the central fibre \mathcal{X}_0 is strictly normal crossings (called such models *snc models*), then the dual intersection complex naturally embeds into X^{an} , its image is called the *skeleton* $\text{Sk}(\mathcal{X})$ associated to the snc model \mathcal{X} . For a holomorphic differential Ω on X Kontsevich and Soibelman have define a so-called *weight function* $\text{wt}_\Omega : X^{\text{an}} \rightarrow \mathbb{R}$. Mustața and Nicaise have studied this function and have shown that it is piecewise affine on any skeleton of a snc model with respect to the natural PL structure. Moreover, Nicaise and Xu have shown that the minimum of wt_Ω is the log canonical threshold. For $x, y \in X^{\text{an}}$ define $x \sim y$ if and only if there exists a path from x to y that intersects the minimality locus of wt_Ω in finitely many points.

Theorem (S. [Sus18a]) The Gromov-Hausdorff limit of $(\mathcal{X}_s, \tilde{\omega}_s)$ the metric graph $\text{Sk}(\mathcal{X}) / \sim$ where \mathcal{X} is any snc model, with the length of an edge of $\text{Sk}(\mathcal{X})$ given by the absolute value of the leading coefficient of the power series expansion of the form Ω in the local coordinate associated to the edge.

The solution uses an approach, due to A. Robinson, to view an algebraically closed valued field as an asymptotic limit of real closed fields; it allows one to treat elements of the field with strictly positive valuation as infinitesimals and to systematically relate the asymptotic behaviour of functions on \mathcal{X}_s as s tends to 0 to non-Archimedean norms of functions on X^{an} . Robinson's asymptotic field has been used in a similar manner by Kramer and Tent [KT02] to compute asymptotic cones of Lie groups.

My next goal in this project is generalization of the theorem above to higher dimensions using the theory of real differential forms of Chambert-Loir and Ducros [CLD12], [Gub16]. This theory defines sheaves of (p, q) -differential forms on X^{an} as objects that are given locally on so-called tropical charts. The latter are the data of an open set $U = \text{trop}^{-1}(\text{trop}(U))$ and a locally closed embedding of U into an algebraic torus. A differential form on a tropical chart U is given by a formal expression involving formal differentials d' and d'' , of bidegree $(1, 0)$ and $(0, 1)$, respectively, of functions $\log |x_1|, \dots, \log |x_n|$ where x_1, \dots, x_n are coordinate functions on the torus.

The support of a positive symmetric $(1, 1)$ form is a polyhedral space and is naturally endowed with a generalized Riemannian structure. I dispose of a tentative definition of the *non-Archimedean limit* of a degenerating family of real differential forms on \mathcal{X}_s with values in the (p, q) -forms on X^{an} which makes precise the remark of Chambert-Loir and Ducros that the forms $d' \log |x|$ and $d'' \log |x|$ in their theory should be regarded as analogues of real forms $d \log_{|s|} |x|$ and $d \text{Arg } x$ on a family of complex manifolds as $s \rightarrow 0$.

The statement that I intend to prove can be formulated as follows:

Claim. Let α be the non-Archimedean limit of Kähler forms $\tilde{\omega}_s$ as $s \rightarrow 0$. Then the Gromov-Hausdorff limit of $(\mathcal{X}_s, \tilde{\omega}_s)$ is isometric (after rescaling so that the diameter is 1) to $\text{supp } \alpha / \sim$ where for all $x, y \in X^{\text{an}}$, $x \sim y$ if and only if there exists a path γ connecting x and y such that $\gamma \cap \text{supp } \alpha$ is finite.

Note that this is a natural generalization of the description of Gromov-Hausdorff limit in the case of flat metrics on curves mentioned above. Establishing this theorem will provide a natural framework for attacking Conjecture C (of course, showing that non-Archimedean limits of Kähler forms of Ricci-flat metrics satisfy the real Monge-Ampere equation remains a considerable challenge). It also provides a tool to compute the collapsed limits of Conjecture A in the odd-dimensional case. Such limits have been studied so far only in the hyperkähler case (see [GW00],[Tos10] [GTZ16], [TZ17]) by reducing them to the so-called large Kähler structure limits, where a complex manifold X is fixed and is assumed to fibre in Lagrangian (with respect to the holomorphic symplectic form) tori over the base of the dimension $1/2 \dim X$, the Kähler form tends towards a boundary of the Kähler cone and the volume of the fibres tends to 0.

2b. Singular affine structure on the essential skeleton.

If X is a variety with trivial canonical class then the essential skeleton $\text{Sk}(X)$ is the minimality locus of the weight function wt_Ω for any $\Omega \in H^0(X, \Omega_X^n)$. Nicaise and Xu [NX16] have shown that

$\text{Sk}(X)$ coincides with the skeleton $\text{Sk}(\mathcal{X})$ of a particular kind of model, a *minimal dlt model*. Although such models are not canonical, $\text{Sk}(\mathcal{X})$ is canonical as a topological space as \mathcal{X} ranges over minimal dlt models. Nicaise, Xu and Yue Yu [NXY18] have found that any minimal dlt model \mathcal{X} defines a surjective map $p_{\mathcal{X}} : X \rightarrow B = \text{Sk}(X)$ which is a fibration in non-Archimedean tori over a subset $B^{\text{sm}} \subset B$ such that $B \setminus B^{\text{sm}}$ is of codimension 2, as predicted by **Conjecture B**. The choice of a model \mathcal{X} corresponds to the choice of a polyhedral decomposition of B and the choice of the fibration $p_{\mathcal{X}}$.

Let $B = \text{Sk}(X)$ and let $p_{\mathcal{X}} : X^{\text{an}} \rightarrow B$ be the retraction map associated to some minimal dlt model. Leaving aside the definition of the pullback $p_{\mathcal{X}}^*$ for the moment, let us formally write $\eta = p_{\mathcal{X}}^* \eta|_B$ for $\eta \in \mathcal{A}^{p,q}(X)$ if the tropical coordinates $-\log|x_1|, \dots, -\log|x_n|$ of all tropical charts used to define η are constant along the fibres of $p_{\mathcal{X}}$. Let α be the non-Archimedean limit of Kähler forms ω_s . If α is a pullback of its restriction to B and is strictly positive on B then it follows from the Claim above (Project 2a) that the Gromov-Hausdorff limit of (\mathcal{X}, ω_s) is homeomorphic to B with the metric structure induced by α .

The Kähler forms of Ricci-flat metrics in a Kähler class $[\omega_0]$ are solutions to the complex Monge-Ampere equation $(\omega_0 + dd^c \varphi)^n = e^f \omega_0^n$ where $f = \log \frac{\Omega \wedge \bar{\Omega}}{\omega_0^n}$ for some holomorphic volume form Ω . The existence of the solution to the non-Archimedean Monge-Ampere equation has been proved by Boucksom, Favre and Jonsson [BFJ15]. We will make the following working assumption:

Assumption. Let α_0 be the non-Archimedean limit of $(\omega_0)_s$, and μ be the non-Archimedean limit of $(\Omega \wedge \bar{\Omega})_s$ as s tends to 0. Let φ_s be the solutions to the complex Monge-Ampere equations above, with the reference Kähler forms $(\omega_0)_s$, for s in a neighbourhood of 0. Then the non-Archimedean limit ψ of φ_s satisfies the non-Archimedean Monge-Ampere equation

$$(\alpha_0 + d'd''\psi)^n = \mu$$

If $\beta = \alpha_0 + d'd''\psi$ then it follows immediately from the definitions of differentials d', d'' that if $p : X^{\text{sm}} \rightarrow B^{\text{sm}}$ is a torus fibration and if $\beta|_{X^{\text{sm}}} = p^*(\beta|_{B^{\text{sm}}})$ then $\beta|_{B^{\text{sm}}}$ is a solution to the real Monge-Ampere equation on B^{sm} in the affine coordinates induced by $p_{\mathcal{X}}$.

Taking the Assumption into account, it is therefore natural to ask the following

Question. Given the first Chern class $c_1(L)$ of the polarization as a non-Archimedean Dolbeault cohomology class of bidegree $(1, 1)$ on X^{an} , find all minimal dlt models \mathcal{X} such that $c_1(L) = p_{\mathcal{X}}^* \beta$ for some $\beta \in H^{1,1}(\text{Sk}(X))$.

2c. Hodge theory for differential forms on non-Archimedean analytic spaces.

As has been already remarked above, a $(1, 1)$ -form in the sense of Chamber-Loir-Ducros defines a generalized Riemannian structure on its support. It comes as no surprise then that it is possible to define a natural L^2 metric on the space of differential forms on X^{an} , and it is natural to ask if there is a non-Archimedean analogue of Kähler condition and Hodge theory.

The notion of a *Kähler superform* on \mathbb{R}^n has been introduced and studied by Lagerberg [Lag11] with certain parts of the complex theory, such as Kähler identities, local dd^c -lemma and Bochner-Kodaira-Nakano identity, carried over to the setting of superforms. The natural Dolbeault complex of (p, q) -forms on non-Archimedean analytic spaces has been studied by Jell, in particular, a version of Poincaré lemma holds for d' (and d'') and therefore $H_{d''}^{p,0}(X^{\text{an}})$ is isomorphic to the p -th singular cohomology group of X^{an} . Furthermore, Jell, Shaw and Smacka [JSS15] have shown that the Dolbeault cohomology groups on a polyhedral space are naturally isomorphic to the tropical cohomology groups defined by Itenberg-Katzarkov-Mikhalkin-Zharkov [IKMZ16].

The theory looks ripe to try to prove non-Archimedean analogues of main statements of Hodge theory on Kähler manifolds and its consequences. In particular, I expect that every Dolbeault cohomology class in $H_{d''}^{p,q}(X)$ has a unique harmonic representative. Furthermore, I expect the global dd^c -lemma, and non-Archimedean versions of Kodaira vanishing and Kodaira embedding theorem to hold. Note that a particular case of the Kodaira embedding theorem is known for non-Archimedean analytic tori, with proofs giving explicit power series representations for the sections of a positive line bundle. There are certain technical peculiarities in the non-Archimedean case, for example, the support of a typical Kähler form is proper closed subset of X^{an} and therefore the theory should be developed under assumption that it contains a skeleton of an snc model (to which X^{an} is homotopy equivalent).

References

- [Art57] Emil Artin. *Geometric algebra*. Interscience, 1957.
- [Bea99] Arnaud Beauville. Counting rational curves on K3 surfaces. *Duke Mathematical Journal*, 97(1):99–108, 1999.
- [BFJ15] Sébastien Boucksom, Charles Favre, and Mattias Jonsson. Solution to a non-Archimedean Monge-Ampere equation. *Journal of the American Mathematical Society*, 28(3):617–667, 2015.
- [BKT10] Fedor Bogomolov, Mikhael Korotiaev, and Yuri Tschinkel. A Torelli theorem for curves over finite fields. *Pure and Applied Mathematics Quarterly*, 6(1):245–294, 2010.
- [CLD12] Antoine Chambert-Loir and Antoine Ducros. Formes différentielles réelles et courants sur les espaces de Berkovich. *arXiv preprint arXiv:1204.6277*, 2012.
- [FGvS99] Barbara Fantechi, Lothar Goettsche, and Duco van Straten. Euler number of the compactified Jacobian and multiplicity of rational curves. *Journal of Algebraic Geometry*, 8:115–133, 1999.
- [GS14] Lothar Göttsche and Vivek Shende. Refined curve counting on complex surfaces. *Geometry & Topology*, 18(4):2245–2307, 2014.
- [GTZ16] Mark Gross, Valentino Tosatti, and Yuguang Zhang. Gromov–Hausdorff collapsing of Calabi–Yau manifolds. *Communications in Analysis and Geometry*, 24(1):93–113, 2016.
- [Gub16] Walter Gubler. Forms and currents on the analytification of an algebraic variety (after Chambert-Loir and Ducros). In *Nonarchimedean and Tropical Geometry*, pages 1–30. Springer, 2016.
- [GW00] Mark Gross and Pelham MH Wilson. Large complex structure limits of K3 surfaces. *Journal of Differential Geometry*, 55(3):475–546, 2000.
- [Har67] Robin Hartshorne. *Foundations of projective geometry*. Harvard University Press, 1967.
- [HK06] Ehud Hrushovski and David Kazhdan. Integration in valued fields. In *Algebraic geometry and number theory*, pages 261–405. Springer, 2006.
- [HS17] Assaf Hasson and Dmitry Sustretov. Incidence systems on Cartesian powers of algebraic curves. *arXiv preprint arxiv:1702.05554*, 2017.
- [IKMZ16] Ilia Itenberg, Ludmil Katzarkov, Grigory Mikhalkin, and Ilia Zharkov. Tropical homology. *Mathematische Annalen*, pages 1–44, 2016.
- [JSS15] Philipp Jell, Kristin Shaw, and Jascha Smacka. Superforms, tropical cohomology, and Poincaré duality. *arXiv preprint arXiv:1512.07409*, 2015.
- [KS04] Maxim Kontsevich and Yan Soibelman. Affine structures and non-Archimedean analytic spaces. *arXiv preprint math/0406564*, 2004.
- [KST11] Martijn Kool, Vivek Shende, and Richard P Thomas. A short proof of the Göttsche conjecture. *Geometry & Topology*, 15(1):397–406, 2011.
- [KT02] Linus Kramer and Katrin Tent. Affine Λ -buildings, ultrapowers of Lie groups and Riemannian symmetric spaces: an algebraic proof of the Margulis conjecture. *arXiv preprint math/0209122*, 2002.
- [Lag11] Aron Lagerberg. L^2 -estimates for the d -operator acting on super forms. *arXiv preprint arXiv:1109.3983*, 2011.
- [NPS16] Johannes Nicaise, Sam Payne, and Franziska Schroeter. Tropical refined curve counting via motivic integration. *arXiv preprint arXiv:1603.08424*, 2016.
- [NX16] Johannes Nicaise and Chenyang Xu. The essential skeleton of a degeneration of algebraic varieties. *American Journal of Mathematics*, 138(6):1645–1667, 2016.
- [NXY18] Johannes Nicaise, Chenyang Xu, and Tony Yue Yu. The non-archimedean SYZ fibration. *arXiv preprint arXiv:1802.00287*, 2018.
- [PT10] Rahul Pandharipande and Richard Thomas. Stable pairs and BPS invariants. *Journal of the American Mathematical Society*, 23(1):267–297, 2010.
- [SSZ14] Vinesh Solanki, Dmitry Sustretov, and Boris Zilber. The quantum harmonic oscillator as a Zariski geometry. *Annals of Pure and Applied Logic*, 165(6):1149–1168, 2014.

- [Sus16] Dmitry Sustretov. Generalised imaginaries and Galois cohomology. *The Journal of Symbolic Logic*, 81(3):917–935, 2016.
- [Sus18a] Dmitry Sustretov. Gromov-Hausdorff limits of flat Riemannian surfaces and non-Archimedean geometry. *arXiv preprint arXiv:1802.03818*, 2018.
- [Sus18b] Dmitry Sustretov. Motivic volume of families of polarized rigid-analytic tori. *arXiv preprint arXiv:1805.04942*, 2018.
- [Tos10] Valentino Tosatti. Adiabatic limits of Ricci-flat Kähler metrics. *Journal of Differential Geometry*, 84(2):427–453, 2010.
- [TZ17] Valentino Tosatti and Yuguang Zhang. Collapsing hyperkähler manifolds. *arXiv preprint arXiv:1705.03299*, 2017.
- [Zil14] Boris Zilber. A curve and its abstract Jacobian. *International Mathematics Research Notices*, 2014(5):1425–1439, 2014.